



TITLE:

# Studies on Holonomic Quantum Fields I (超函数と線型微分方程式 V)

AUTHOR(S):

SATO, MIKIO; MIWA, TETSUJI; JIMBO, MICHIO

---

CITATION:

SATO, MIKIO ...[et al]. Studies on Holonomic Quantum Fields I (超函数と線型微分方程式 V). 数理解析研究所講究録 1977, 287: 42-48

ISSUE DATE:

1977-02

URL:

<http://hdl.handle.net/2433/106132>

RIGHT:

# Studies on Holonomic Quantum Fields. I

By

Mikio SATO, Tetsuji MIWA and Michio JIMBO

To understanding the mathematical structure of quantized fields or systems with infinite freedom, non trivial but exactly calculable models would be of great help [1]. In this and subsequent notes we present, both in the continuum and in the lattice, 2-dimensional soluble models of neutral scalar massive field theory whose  $\tau$ -functions exhibit a non trivial singularity structure.

In the present article we deal with the continuum case. We introduce an auxiliary free fermi/bose field and construct the field operator by giving its induced rotation in the space of wave functions. Making use of the "theory of rotation" (2. cf.[2]) developed recently by the first author, we express this field operator in the normal product form of these free fields. We also calculate the asymptotic fields and the S-matrix of the field  $\varphi^F$  defined in 3. Next we give explicit formulae for  $\tau$ -functions of these models and study their holonomy structure.

The lattice field theory will be discussed in a subsequent paper. Specifically we shall show that our model  $\varphi^F / \varphi_F$  coincide with the scaling limit of the Ising model from above/below the critical temperature. Main part of these results has been announced in [3].

We use the following notations. The space-time and the energy-momentum co-ordinates are denoted by  $x=(x^0, x^1)$  and

$p=(p^0, p^1)$ . We also use  $x^\pm = (x^0 \pm x^1)/2$  and  $p^\pm = p^0 \pm p^1$ . The mass-shell  $\{p \in \mathbb{R}^2 | p^2 = (p^0)^2 - (p^1)^2 = m^2\}$  ( $m > 0$ ) is denoted by  $M$ . For  $p \in M$  we set  $u^{\pm 1} = p^\pm / m$ ,  $du = du/2\pi |u|$ .

1. Let  $\psi(u)^\dagger$  and  $\psi(u)$  ( $u > 0$ ) be the creation and annihilation operators of auxiliary fermion. If we define  $\psi(u) = \psi(-u')^\dagger$  for  $u < 0$ , their anti-commutation relation reads  $[\psi(u), \psi(u')]_+ = 2\pi |u| \delta(u+u')$ . Likewise we define auxiliary bosons  $\phi(u)$  with the commutation relation  $[\phi(u), \phi(u')]_- = 2\pi u \delta(u+u')$ . In two dimensional space-time these two are in fact equivalent. Namely

$$(1) \quad \phi_\pm(u) = : \psi(u) \exp \int_0^\infty \theta(\pm(|u| - u')) \psi(u')^\dagger \psi(u') du' :$$

satisfy the commutation relation  $[\phi_\pm(u), \phi_\pm(u')]_- = 2\pi u \delta(u+u')$ , and conversely  $\psi(u)$  is given by

$$(2) \quad \psi(u) = : \phi_\pm(u) \exp \int_0^\infty \theta(\pm(|u| - u')) \phi_\pm(u')^\dagger \phi_\pm(u') du' : .$$

2. We let  $W$  denote an orthogonal/symplectic space, a vector space equipped with a non-degenerate symmetric/skew-symmetric inner product  $\langle w, w' \rangle$ . First consider the orthogonal case and denote by  $A(W)$  the enveloping algebra (Clifford algebra) over  $W$  with defining relation  $[w, w']_+ = \langle w, w' \rangle$ .  $G(W)$  denotes the Clifford group  $\{g \in A(W) | \exists g^{-1}, gwg^{-1} = w\}$ . Let  $g \mapsto g^*$  denote the anti-automorphism of  $A(W)$  characterized by  $w^* = w$  for  $w \in W$ . Set  $n(g) = g^*g = gg^*$  for  $g \in G(W)$ , and  $g \mapsto n(g)$  defines a group homomorphism  $G(W) \rightarrow GL(1)$ . Let  $W = V^\dagger \oplus V$  be a decomposition into two holonomic subspaces. This means that there exist a basis  $\psi^\dagger = (\psi_\mu^\dagger)$  of  $V^\dagger$  and a basis  $\psi = (\psi_\mu)$  of  $V$  such that  $\langle \psi_\mu^\dagger, \psi_\nu^\dagger \rangle = 0$ ,  $\langle \psi_\mu, \psi_\nu \rangle$

$=0$  and  $\langle \psi_\mu^\dagger, \psi_\nu \rangle = \delta_{\mu\nu}$ . Then  $A(W)$  is a semi-direct product of two exterior algebras  $\Lambda(V^\dagger)$  and  $\Lambda(V)$ , and a  $\Lambda(V^\dagger) \rightarrow \Lambda(V)$ -isomorphism  $N: A(W) = \Lambda(V^\dagger) \cdot \Lambda(V) \rightarrow \Lambda(W) = \Lambda(V^\dagger) \wedge \Lambda(V)$  such that  $N(1) = 1$  is determined uniquely. The image  $N(g) \in \Lambda(W)$  we call the norm of  $g$ . (In physicist's notation  $g =: N(g) :.$ ) For  $g \in G(W)$   $T_g: w \in W \mapsto gw g^{-1} \in W$  is a rotation, an isomorphism which preserves the inner product. Let  $T_g(\psi^\dagger, \psi) = (\psi^\dagger, \psi) \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix}$ . First assume that  $T_4$  is invertible. Then we have the following expression of the norm of  $g$ .

$$(3) \quad N(g) = \langle g \rangle \exp((1/2) \psi^\dagger T_2 T_4^{-1} \psi + \psi^\dagger (T_4^{-1} - 1) \psi + (1/2) \psi T_4^{-1} T_3 \psi),$$

where  $n(g) = \langle g \rangle^2 (\det T_4)^{-1}$ , and we regard  $\psi_\mu^\dagger, \psi_\mu$  as elements of  $\Lambda(W)$ . Next we assume that  $\dim \text{Ker } T_4 = 1$ , and choose  $\psi_0^\dagger \in V^\dagger$ ,  $\psi_0 \in V$  and  $w \in G(W) \cap W$  such that  $T_g \psi_0 = \psi_0^\dagger$ ,  $w^2 = 1$  and  $\langle w, \psi_0^\dagger \rangle = 1$ . Then  $(T_{wg})_4$  is invertible and

$$(4) \quad N(g) = \psi_0^\dagger N(wg) + N(wg) \psi_0.$$

Here we regard  $\psi_0^\dagger$  and  $\psi_0$  as elements of  $\Lambda(W)$ . Next consider the symplectic case, and define  $A(W)$ ,  $G(W)$ , etc. with due modifications. In particular  $w^* = iw$  for  $w \in W$ , and the norm of  $g \in A(W)$  is defined as an element of the symmetric tensor algebra  $S(W)$ . Assuming that  $T_4$  is invertible, we have

$$(5) \quad N(g) = \langle g \rangle \exp((1/2) \phi^\dagger (-T_2 T_4^{-1}) \phi + \phi^\dagger (T_4^{-1} - 1) \phi + (1/2) \phi T_4^{-1} T_3 \phi),$$

with  $n(g) = \langle g \rangle^2 \det T_4$ .

3. Let now  $W$  be the space of wave functions  $w(x) = (w_+(x), w_-(x))$  satisfying the Dirac equation  $\partial w_\pm(x) / \partial x^\pm \mp m w_\mp(x) = 0$ . An orthogonal structure is introduced to  $W$  by

defining  $\langle w, w' \rangle = \int_{-\infty}^{+\infty} dx^1 (w_+(x) w'_+(x) + w_-(x) w'_-(x))$ . If we identify  $w \in W$  with the operator  $\int_{-\infty}^{+\infty} dx^1 (w_+(x) \psi_+(x) + w_-(x) \psi_-(x))$ , where  $\psi_{\pm}(x) = (1/\sqrt{2}) \int_{-\infty}^{+\infty} \frac{du}{\sqrt{0+iu}} \psi(u) \exp(-im(x^-u + x^+u^{-1}))$ , the Clifford algebra  $A(W)$  is nothing but the operator algebra of free fermions. We choose as  $V^\dagger/V$  the set of creation/annihilation operators in  $W$ . Set  $W_x^+ = \{w \in W | w(x') = 0 \text{ if } (x' - x)^2 < 0, x'^1 - x^1 \leq 0\}$ , and we shall have  $W = W_x^+ \oplus W_x^-$ , an orthogonal decomposition. We now introduce our field operator  $\varphi_F(x) \in A(W)$  by specifying its induced rotation  $T_{\varphi_F(x)}$  with the property  $T_{\varphi_F(x)}^2 = 1$  by

$$(6) \quad T_{\varphi_F(x)}(w^+ + w^-) = w^+ - w^-, \quad w^\pm \in W_x^\pm.$$

Applying the formula (3) to the present situation and choosing  $\langle \varphi_F(x) \rangle = 1$  we obtain the following expression for  $\varphi_F(x)$ :

$$(7) \quad \varphi_F(x) = : \exp L_F(x) :,$$

where  $L_F(x) = (1/2) \int \int_{-\infty}^{+\infty} \frac{du}{\sqrt{0+iu}} \frac{du'}{\sqrt{0+iu'}} \frac{-1(u-u')}{u+u'-i0} \psi(u) \psi(u') \exp(-im(x^-(u+u') + x^+(u^{-1}+u'^{-1})))$ . The micro-causality and the Lorentz covariance of  $\varphi_F(x)$  are manifest in this approach.

We construct  $\varphi^F(x)$  and  $\varphi_B(x)$  analogously, using the formulae in the case  $\dim \text{Ker } T_4 = 1$  and in the symplectic case, respectively. In the latter case we choose as  $W$  the solution space to the Klein-Gordon equation and equip it with the inner product  $\langle w, w' \rangle = -i \int_{-\infty}^{+\infty} dx^1 (w(x) \cdot \partial w'(x) / \partial x^0 - w(x) / \partial x^0 \cdot w'(x))$ . The results are

$$(8) \quad \varphi^F(x) = : \psi_0(x) \exp L_F(x) :,$$

where  $\psi_0(x) = \int_{-\infty}^{+\infty} du \psi(u) \exp(-im(x^-u + x^+u^{-1}))$ ,

$$(9) \quad \varphi_B(x) = : \exp L_B(x) :,$$

where  $L_B(x) = (1/2) \iint_{-\infty}^{+\infty} \frac{du}{u} \frac{du'}{u'} \frac{-2\sqrt{u-i0}\sqrt{u'-i0}}{u+u'-i0} \phi(u)\phi(u')$   
 $\exp(-im(x^-(u+u') + x^+(u^{-1}+u'^{-1})))$ .

4. The asymptotic fields for  $\varphi^F$  are defined by

$$(10) \quad \varphi_{\pm}(x) = \int_{-\infty}^{+\infty} \frac{du}{u} \varphi_{\pm}(u) \exp(-ipx),$$

where  $\varphi_{\pm}(u) = \lim_{t \rightarrow \pm\infty} i \int_{x^0=t} dx^1 (\varphi^F(x) (\partial/\partial x^0) \exp(ipx) - (\partial/\partial x^0) \varphi^F(x) \exp(ipx))$ . We find that this limit coincides with  $\phi_{\pm}(u)$  defined in 1. The asymptotic states  $|>_{\pm}$  are related to the auxiliary fermion states  $|>_F$  through the formulae

$$(11) \quad |u_n, \dots, u_1>_{\pm} = \prod_{i < j} \varepsilon(\pm(u_i - u_j)) |u_n, \dots, u_1>_F,$$

where  $\varepsilon(u)$  stands for the signature of  $u$ . Accordingly the particle number is conserved, and the S-matrix in the n-particle state is given by  $(-)^{n(n-1)/2}$  times the identity operator, showing that the maximum phase shift is attained in this model.

5. The n-point  $\tau$ -function of an operator  $\varphi(x)$  is expressed as follows:

$$(12) \quad \tau_n(p_1, \dots, p_n) = \sum_{\text{permutations}} \frac{n!}{n-1} T_{n-1}(p_1, p_1+p_2, \dots, p_1+\dots+p_{n-1})$$

$$\times (2\pi)^2 \delta^2(p_1+\dots+p_n),$$

$$T_{n-1}(q_1, \dots, q_{n-1}) = \sum_v (1/v_1! \dots v_{n-1}!) \int_0^{\infty} \frac{du}{u} \prod_{j=1}^n \varphi_{v_j+v_{j-1}}(u_{jv_j}, \dots, u_{jv_j} - u_{j-1v_{j-1}}, \dots, -u_{j-1v_{j-1}})$$

$$\times \prod_{j=1}^{n-1} 2\pi \delta(q_j^+ - mU_j) i(q_j^- - mU_j' + i0)^{-1},$$

with  $U_j = \sum_{k=1}^{v_j} u_{jk}$ ,  $U_j' = \sum_{k=1}^{v_j} u_{jk}^{-1}$  and  $v_0 = v_n = 0$ . The (anti-)

symmetric functions  $\varphi_n$  are the matrix elements defined by  $\varphi_n(u_1, \dots, u_n) = \langle -u_{m+1}, \dots, -u_n | \varphi(0) | u_1, \dots, u_m \rangle$  for  $u_1, \dots, u_m > 0$  and  $u_{m+1}, \dots, u_n < 0$ . In our models they are obtained from (7), (8) and (9).

$$(13) \quad \varphi_{F,n}(u_1, \dots, u_n) = \text{Pfaffian} \left( i P \frac{u_j - u_k}{u_j + u_k} \right)_{1 \leq j, k \leq n}$$

$$= \begin{cases} i^{n/2} \prod_{1 \leq j < k \leq n} P \frac{u_j - u_k}{u_j + u_k} & (n \text{ even}) \\ 0 & (n \text{ odd}), \end{cases}$$

$$(14) \quad \varphi_n^F(u_1, \dots, u_n) = -i \varphi_{F,n+1}(\infty, u_1, \dots, u_n)$$

$$= \begin{cases} 0 & (n \text{ even}) \\ i^{(n-1)/2} \prod_{1 \leq j < k \leq n} P \frac{u_j - u_k}{u_j + u_k} & (n \text{ odd}), \end{cases}$$

and

$$(15) \quad \varphi_{B,n}(u_1, \dots, u_n) = \text{Hafnian} \left( -2 P \frac{\sqrt{u_j - i0} \sqrt{u_k - i0}}{u_j + u_k} \right)_{1 \leq j, k \leq n}.$$

Here  $P(1/u+v)$  denotes the principal value of  $1/u+v$ , and for a symmetric matrix  $(a_{jk})_{1 \leq j, k \leq n}$  we set  $\text{Hafnian}(a_{jk}) = 0$  for odd  $n$  and  $= \sum a_{j_1 j_2} a_{j_3 j_4} \dots a_{j_{n-1} j_n}$  for even  $n$ , where the sum is taken over  $(n-1)!!$  pairings of indices  $1, \dots, n$ . In particular the (Euclidean) two point functions of  $\varphi_F$  and  $\varphi^F$  coincide with those obtained by [4] and [5].

The singularity/holonomy spectrum of  $\tau_n(p)$  is confined to the union of positive- $\alpha$ /complex Landau singularities

corresponding to graphs with no internal vertices [6], where the number of (internal and external) lines incident to each vertex is always even for  $\varphi^F$  and is always odd for  $\varphi_F$ ,  $\varphi_B$ . On the leading singularity  $\Lambda_G^+$ , the order of  $\tau_n$  for  $\varphi^F$  or  $\varphi_F$  is given by

$$(16) \quad \text{ord}_{\Lambda_G^+} \tau_n = n_e - N/2 - \sum_{i < j} N_{ij} (N_{ij} - 1)/2,$$

where  $n_e$  denotes the number of vertices of  $G$ ,  $N_{ij}$  the number of internal lines joining the vertices  $i$  and  $j$ , and  $N = \sum_{i < j} N_{ij}$ . Note that repulsive effect of multiple internal lines is incorporated in (16).

Finally we remark that the generalized unitarity relation for the  $\tau$ -function of  $\varphi^F$

$$(17) \quad 0 = \sum_{\ell=0}^{\infty} \frac{1}{\ell!} \sum_{k=0}^n (-)^k \sum_{\text{combinations}} \binom{n}{k} \int \cdots \int \prod_{i=1}^{\ell} du_i \tau_k^{(\ell)}(p_1, \dots, p_k; u_1, \dots, u_{\ell})$$

$$\times \overline{\tau_{n-k}^{(\ell)}(-p_{k+1}, \dots, -p_n; u_1, \dots, u_{\ell})}$$

where we set  $\tau_k^{(\ell)}(p_1, \dots, p_k; u_1, \dots, u_{\ell}) = \tau_{k+\ell}(p_1, \dots, p_k, q_1, \dots, q_{\ell})$   
 $\times \prod_{i=1}^{\ell} (q_i^2 - m^2) |_{q_i^{\pm} \mapsto u_i^{\pm}}$  and bar denotes the complex conjugation, is

directly and analytically verified by using our explicit formulae (12) and (14).

#### References

- [1] M. Sato: in Proc. MΦ, Kyoto, 1975. Springer Lecture Notes in Phys. 39.
- [2] B. Kaufman: Phys. Rev. 76, 1232-1243(1949).
- [3] M. Sato, T. Miwa and M. Jimbo: Field theory of the 2-dimensional Ising model in the scaling limit, RIMS preprint 207, (1976).
- [4] T. T. Wu, B. M. McCoy, C. A. Tracy and E. Barouch: Phys. Rev. B 13, 316-374 (1976).
- [5] R. Z. Bariev: Phys. Lett. 55A, 456-458 (1976).
- [6] M. Sato, T. Miwa, M. Jimbo and T. Oshima: Holonomy structure of Landau singularities and Feynman integrals, to appear in Publ. RIMS 12. suppl.